

## PATHWAYS ADDITIONAL REFERENCES NOT IN MAGAZINE: Spring 2016

### Research Perspective

### Symptom Reporting as a Useful Tool to Detect Lymphedema

By Mei R. Fu

#### References

1. Paskett ED, Naughton MJ, McCoy TP, Case LD, Abbott JM. The epidemiology of arm and hand swelling in premenopausal breast cancer survivors. *Cancer Epidemiol Biomarkers Prev.* 2007; 16: 775-782 [PMID: 17416770 DOI: 10.1158/1055-9965.epi-06-0168]
2. McLaughlin SA, Bagaria S, Gibson T et al. Trends in risk reduction practices for the prevention of lymphedema in the first 12 months after breast cancer surgery, *J Am Coll Surg.* 2013; 216:380-389.
3. Fu MR & Kang Y. Psychosocial impact of living with cancer-related lymphedema. *Semin Oncol Nurs.* 2013; 29(1), 50-60.
4. Fu MR, Ridner S, Hu SH et al. Psychosocial Impact of Lymphedema: A systematic review of literature from 2004 to 2011. *Psychooncology.* 2013; 22(7):1466-84. doi: 10.1002/pon.3201. Epub 2012
5. Fu MR, Rosedale M. Breast cancer survivors' experiences of lymphedema-related symptoms. *J Pain Symptom Manag.* 2009; 38:849-59.
6. Chachaj A, Malyszczak K, Pyszel K et al. Physical and psychological impairments of women with upper lymphedema following breast cancer treatment. *Psychooncology.* 2009; 19: 299-305.
7. Cormier JN, Xing Y, Zaniletti I et al. Minimal limb volume change has a significant impact on breast cancer survivors. *Lymphology.* 2009; 42:161-175.
8. Fleissig A, Fallowfield LJ, Langridge CI, Johnson L, Newcombe RG, Dixon JM, et al. Post-operative arm morbidity and quality of life. Results of the ALMANAC randomized trial comparing sentinel node biopsy with standard axillary treatment in the management of patients with early breast cancer. *Breast Cancer Research & Treatment.* 2006; 95: 279-293.
9. Armer JM, Radina ME, Porock D et al. (2003). Predicting breast cancer-related lymphedema using self-reported symptoms. *Nurs Res.*2003; 52(6), 370-379.
10. International Society of Lymphology. The diagnosis and treatment of peripheral lymphedema. *Lymphology.*2003; 36, 84-91.

11. McLaughlin SA, Wright MJ, Morris KT, Giron GL, Sampson MR, Brockway JP, Van Zee KJ. Prevalence of lymphedema in women with breast cancer 5 years after sentinel lymph node biopsy or axillary dissection: objective measurements. *J Clin Oncol.* 2008; 26(32), 5213-5219. doi: 10.1200/JCO.2008.16.3725
12. Hu S, Cleland CM, Kang Y, Fu MR. Measuring lymphedema symptom burdens: A psychometric study. The Multinational Association of Supportive Care in Cancer's annual meeting (MASCC/ISOO): 2012 International Symposium on Supportive Care in Cancer in New York City on June 28-30, 2012.
13. Shi S, Lu Q, Fu MR et al. Psychometric properties of the breast cancer and lymphedema symptom experience index: the Chinese version. *Eur J Oncol Nurs.* pii: S1462-3889(15)00076-9. doi: 10.1016/j.ejon.2015.05.002. [Epub ahead of print] PMID: 26071198
14. Fu MR, Chen C, Haber J et al. The effect of providing information about lymphedema on the cognitive and symptom outcomes of breast cancer survivors. *Ann Surg Oncol.* 2010 Jul; 17(7):1847-53. doi: 10.1245/s10434-010-0941-3. Epub 2010 Feb 6. PMID: 20140528.
15. Youden W J. (1950) Index for rating diagnostic tests. *Cancer*, 3, 32–35.
16. Perkins NJ & Schisterman EF. The Inconsistency of "Optimal" Cutpoints Obtained using Two Criteria based on the Receiver Operating Characteristic Curve. *American J Epidemiol.* 2006; 163(7), 670–675.
17. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria. 2013; ISBN 3-900051-07-0, URL <http://www.R-project.org/>.
18. Smoot BJ, Wong JF, Dodd MJ. Comparison of diagnostic accuracy of clinical measures of breast cancer related lymphedema: Area under the curve. *Arch Phys Med Rehabil* 2011; 92:603-10.
19. Welch BL. On the comparison of several mean values: an alternative approach. *Biometrika*, 1951; 38, 330–336.
20. Fu MR, Axelrod D, Guth, A, Cartwright-Alcarese F, Qiu Z, Goldberg J, Kim J, Scagliola J, Kleinman R, Haber J. Proactive approach to lymphedema risk reduction: a prospective study. *Ann Surg Oncol* 2014; 21(11), 3481-3498. Online First. DOI: 10.1245/s10434-014-3761-z.